

# UNCLASSIFIED

AD NUMBER
AD815244
NEW LIMITATION CHANGE
TO Approved for public release, distribution unlimited
FROM Distribution authorized to U.S. Gov't. agencies and their contractors; Critical Technology; DEC 1965. Other requests shall be referred to Army Biological Laboratories, Fort Detrick, Frederick, MD.
AUTHORITY
SMUFD D/A ltr, 15 Feb 1972

THIS PAGE IS UNCLASSIFIED

INFLUENCE OF THE COMPOSITION OF THE NUTRIENT MEDIUM ON THE COURSE OF LYSIS  
OF PLAGUE CULTURES UNDER THE INFLUENCE OF PLAGUE BACTERIOPHAGE

AD815244

Translation No. 1721

DECEMBER 1965

DDC  
RECEIVED  
JUN 14 1967  
RECEIVED  
C

STATUS: UNCLASSIFIED

This document is subject to export controls and each  
transmission of this document or foreign nationals may be  
made only with prior approval of U. S. ARMY

BIOLOGICAL LABORATORIES  
FORT DETRICK, FREDERICK, MARYLAND

DDC AVAILABILITY NOTICE

Qualified requestors may obtain copies of this document from DDC.

This publication has been translated from the open literature and is available to the general public. Non-DOD agencies may purchase this publication from Clearinghouse for Federal Scientific and Technical Information, U. S. Department of Commerce, Springfield, Va.

Technical Library Branch  
Technical Information Division

INFLUENCE OF THE COMPOSITION OF THE NUTRIENT MEDIUM ON THE COURSE OF LYSIS  
OF PLAGUE CULTURES UNDER THE INFLUENCE OF PLAGUE BACTERIOPHAGE

Following is the translation of an article by T. P. Kudinova, published in the Russian-language periodical Materialy Nauchnoy Konferentsii po Prirodnoy Ochagovosti i Profilaktike Chumy (Materials from the Scientific Conference on the Natural Focality and Prophylaxis of Plague) Alma-Ata, Feb., 1963, pages 129--130. Translation performed by Sp/7 Charles T. Ostertag, Jr.

A study was made on the course of lysis of a plague microbe culture under the influence of plague bacteriophage in a medium, poor in nutrient substances.

Into a physiological solution we added a culture of the plague microbe (EV strain), plague bacteriophage and Hottinger broth up to a content of it in the medium of 1:25 (with a calculation of the broth, introduced with the bacteriophage). As a control we took pure broth with the same amount of bacteriophage and microbial cells. After specific intervals of time a determination was made of the content of viable microbial cells in a unit of volume of the medium (by the method of inoculation on agar following preliminary neutralization of the bacteriophage with the neutral red dye) and the titer of bacteriophage was determined.

After one day following inoculation it was possible to isolate individual microbial cells from the broth with the plague bacteriophage only in rare cases. In the medium consisting of the physiological solution with the addition of 1:25 broth to it, the number of viable cells increased steadily, and after five--six days exceeded their initial content in the medium by 10--15 and even several dozen times, just as lysis of a plague culture takes place under the influence of pseudotuberculosis bacteriophage.

At the same time the increase in the titer of bacteriophage in a physiological solution with broth did not differ essentially from the same in broth, and also remained the same or exceeded by one dilution when titrated according to the method of Appelman.